## Amendments to the Claims:

Please delete all prior listings of claims and substitute therefor the following listing of claims:

## Claims 1-26 (Cancelled)

١

27. (previously presented) A method for obtaining an isolated polynucleotide comprising a sequence encoding a protein having Rubisco carboxylation activity, the method comprising:

recombining a plurality of parental polynucleotide species encoding at least one protein having Rubisco carboxylation activity under conditions suitable for sequence shuffling to form a resultant library of sequence-shuffled polynucleotides;

transferring said library into a plurality of host cells, thereby forming a library of transformants wherein sequence-shuffled Rubisco polynucleotides are expressed;

identifying at least one transformant from said library that expresses an enhanced protein having a Rubisco carboxylation activity that is enhanced to an extent that is statistically significant relative to the Rubisco carboxylation activity of proteins encoded by the plurality of parental polynucleotide species, wherein the identified transformant contains a polynucleotide comprising a sequence encoding the enhanced protein; thereby obtaining a polynucleotide comprising a sequence encoding the enhanced protein.

- 31. (currently amended) The method of claim 27, wherein the enhanced protein has a Km for CO<sub>2</sub> that is less than that of proteins encoded by the plurality of <u>parental</u> polynucleotide species, to an extent that is statistically significant.
- 32. (currently amended) The method of claim 27, wherein the enhanced protein has a Km for CO<sub>2</sub> that is greater than that of proteins encoded by the plurality of <u>parental</u> polynucleotide species, to an extent that is statistically significant.
- 33. (previously presented) The method of claim 27, wherein the plurality of parental polynucleotide species encodes at least one Rubisco Form I L subunit.

- 34. (previously presented) The method of claim 27, wherein the plurality of parental polynucleotide species encodes at least one Rubisco Form I S subunit.
- 35. (previously presented) The method of claim 27, wherein the plurality of parental polynucleotide species encodes at least one Rubisco Form II subunit.
- 36. (previously presented) The method of claim 27 further comprising a selectable marker gene which affords a means of selection when expressed in chloroplasts.

ij.

ì

37. (previously presented) The method of claim 36, wherein the sequence encoding the enhanced protein and the selectable marker gene are flanked by an upstream flanking recombinogenic sequence having sufficient sequence identify to a chloroplast genome sequence to mediate efficient recombination and a downstream flanking recombinogenic sequence having sufficient sequence identify to a chloroplast genome sequence to mediate efficient recombination.

Respectfully submitted,

McANDREWS, HELD & MALLOY, LTD.

By:

Donald J. Pochopien Registration No. 32,167 Attorney for Applicants 500 West Madison Street, 34<sup>th</sup> Floor

Chicago, Illinois 60661

(312) 775-8133

Date: January 10, 2005 J:\open/DJP\Codexis\USPTO\16184US03\Listing of Claims.doc